

[illegible]

D—D°Ñ,DµD¼ Ñ•D¾Ñ,D¿Ñ€D°D²D,D»D°Ñ•ÑƒE D½D° Ñ•D°D¼Ñf D¿Ñ€D¾Ñ†DµD´ÑfÑ€Ñf.
DœDµD½Ñ•D¿D¾D¿Ñ€D¾Ñ•D,D»D,D½D°D´DµÑ,ÑƒE ÑD°D»D°Ñ,D,D»DµÑ‡ÑƒE D½D°
D°ÑfÑ´DµÑ,D°Ñf,D½D° D»DµD²Ñ‹D¹ D±D¾D°. D“ÑfD±Ñ‹D¼D½Dµ Ñ•D¼D°D•D°D»D,
ÑfD²D»D°D¶D½Ñ•ÑŽÑ%œ,D¼ Ñ•Ñ€DµD´Ñ•Ñ,D²D¾D¼,D¿D¾Ñ•D»Dµ Ñ‡DµD³D¾ D²Ñ•
Ñ,D°D²D,D»D,D•D°D³ÑfD±D½D,D°. D´D²DµD½Ñf D²Ñ•Ñ,D°D²D,D»D,D°D°Ñ,DµÑ,DµÑ€D,
D¿D¾D´D°D»ÑŽÑ†D,D»D,D,D•D¼DµÑ€DµD½D,DµD´D°D²D»DµD½D,Ñ•D,D¿ÑfD»ÑƒEÑ•
D°D´D²DµD»D,D¿DµÑ€D²ÑfÑŽ D¿Ñ€D¾D±D½ÑfÑŽ D´D¾D•Ñf D°D½DµÑ•Ñ,DµD•D,D,D´D»Ñ•
D²Ñ‹Ñ•D²D»DµD½D,Ñ•D²D¾D•D¼D¾D¶D½Ñ‹Ñ D¿D¾D±D¾Ñ‡D½Ñ‹Ñ Ñ€DµD°D°Ñ†D,D¹.
D“D¾D»D¾D²D°D•D°D°Ñ€ÑfD¶D,D»D°Ñ•ÑƒE.D´D³D¾Ñ€D»D¾D²D¿Ñ€Ñ‹Ñ•D½ÑfD»D,Ñ€D°Ñ•
Ñ,D²D¾Ñ€D»DµD´D¾D°D°D,D½D°D,D½D°Ñ‡D°D»D,D²D²DµD´DµD½D,DµD¾Ñ•
D½D¾D²D½D¾D¹ D´D¾D•Ñ‹D¿Ñ€DµD¿D°Ñ€D°Ñ,D°.

DŽŇ, D°D»ŇŽŇ†D, D»D°Ň•NCE Ň• DĠŇŇD°D°Ň, D, Ň†DμŇ•D°D, D¼D³D¹½D³¼D²DμD¹½D¹½D³¼. DŸŇŇD³¼Ň
 •D¹½ŇfD»D°Ň•NCE Ň•D°D¼D³¼Ň•Ň, D³¼Ň•Ň, DμD»ŇŇD¹½D³¼, Ň†DμŇŇDμD. 40 D¼D, D¹½ŇfŇ, DŸŇŇD, Ň•Ň, D³¼D¼ D»DμDŲD°D»D° ŇfDŲDμ D¹½D° Ň•DĠD, D¹½Dμ, D±DμD. D°D°Ň, DμŇ, DμŇŇD° D,
 D•D°D³ŇfD±D¹½D, D°D°, DĠD³¼D´ D³¼D´DμŇ•D»D³¼D¼. DœDμD¹½Ň•
 DĠŇŇD³¼D, D¹½Ň, D³¼ŇŇD¼D, ŇŇD³¼D²D°D»D, Ň†Ň, D³¼ D³D°Ň•Ň, ŇŇD³¼Ň•D°D³¼DĠD, Ň•
 D´D»D, D»D°Ň•NCE 7 D¼D, D¹½ŇfŇ, DƒD³¼Ň•D¹½D³¼Ň, Ň< D, D³¼Ň%ŇfŇ% DμD¹½D, D¹ D²
 DŲDμD»ŇfD´D°Dμ D¹½Dμ D¹½D°D±D»ŇŽD´D°D»D³¼Ň•NCE. DŠD°D•D°D»D³¼Ň•NCE, Ň• DĠŇŇD³¼Ň•
 DĠD°D»D° D¼D°D°Ň•D, D¼ŇŇfD¼ DĠD³¼D»D¼D, D¹½ŇfŇ, Ň<, D¹½D³¼ DĠŇŇD, Ň•Ň, D³¼D¼ Ň†ŇfD²Ň•
 Ň, D²D³¼D²D°D»D° Ň•DμD±Ň• D³¼Ň, D´D³¼Ň D¹½ŇfD²Ň•DμD¹. DœDμD¹½Ň• DĠD³¼DĠŇŇD³¼Ň•D, D»D,
 DĠŇŇD, Ň•DμŇ•Ň, ŇŇD D¹½D° D°ŇfŇ•DμŇ, D°Dμ D, DĠD³¼Ň•D, D´DμŇ, ŇŇD DĠD°ŇŇŇf D¼D, D¹½ŇfŇ, ,
 DĠD³¼Ň•D»Dμ Ň†DμD³D³¼ DĠD³¼D¼D³¼D³D»D, D¼D¹½Dμ D²Ň•Ň, D°Ň, ŇŇD. DƒD±DμD´D, D²Ň•D, Ň•
 ŇŇD, Ň†Ň, D³¼ Ň•Ň, D³¼ŇŽ ŇfD²DμŇŇDμD¹½D¹½D³¼, DĠD³¼DĠŇŇD³¼Ň•D, D»D, DĠŇŇD³¼D¹Ň, D, D²
 D°D³¼ŇŇD, D´D³¼ŇŇD, D³¼DŲD, D´D°Ň, ŇŇD ŇŇDμD•ŇfD»ŇŇŇ, D°Ň, D° D³D°Ň•Ň, ŇŇD³¼Ň•
 D°D³¼DĠD, D, .

$\partial \tilde{Y} \tilde{N} \in \mathcal{D}, \mathcal{D}^{1/4} \mathcal{D} \mu \tilde{N} \in \mathcal{D}^{1/2} \mathcal{D}^{3/4} \tilde{N} \nmid \mathcal{D} \mu \tilde{N} \in \mathcal{D} \mu \mathcal{D} \cdot 10 \mathcal{D}^{1/4} \mathcal{D}, \mathcal{D}^{1/2} \tilde{N} f \tilde{N}, \mathcal{D}^2 \tilde{N} \in \mathcal{D}^\circ \tilde{N} \nmid \mathcal{D} \zeta \mathcal{D}^{3/4} \mathcal{D} \cdot \mathcal{D}^2 \mathcal{D}^\circ \mathcal{D} \gg \mathcal{D}^\circ \mathcal{D}^{1/4} \mathcal{D} \mu \mathcal{D}^{1/2} \tilde{N} \cdot$
 $\mathcal{D}, \tilde{N} \cdot \mathcal{D}^{3/4} \mathcal{D}^{3/4} \mathcal{D} \pm \tilde{N} \% \circ \mathcal{D}, \mathcal{D} \gg \mathcal{D}^\circ \tilde{N} \in \mathcal{D} \mu \mathcal{D} \cdot \tilde{N} f \mathcal{D} \gg \tilde{N} \mathcal{C} \tilde{N}, \mathcal{D}^\circ \tilde{N}, \tilde{N} \prec \mathcal{D}, \mathcal{D}^2 \tilde{N} \prec \mathcal{D}' \mathcal{D}^\circ \mathcal{D} \gg \mathcal{D}^\circ \mathcal{D}^{1/4} \mathcal{D}^{1/2} \mathcal{D} \mu$
 $\tilde{N} \nmid \mathcal{D}^2 \mathcal{D} \mu \tilde{N}, \mathcal{D}^{1/2} \tilde{N} f \tilde{N} \tilde{Z} \tilde{N} \in \mathcal{D}^\circ \tilde{N} \cdot \mathcal{D} \zeta \mathcal{D} \mu \tilde{N} \nmid \mathcal{D}^\circ \tilde{N}, \mathcal{D}^\circ \tilde{N} f \tilde{N} \cdot \tilde{N} \mathcal{D}^\circ \tilde{N} \in \mathcal{D}^\circ \mathcal{D}^\circ \tilde{N}, \mathcal{D} \mu \tilde{N} \in \mathcal{D}^{1/2} \tilde{N} \prec \mathcal{D}^{1/4} \mathcal{D},$
 $\mathcal{D}^{1/4} \mathcal{D}^{3/4} \mathcal{D}^{1/4} \mathcal{D} \mu \mathcal{D}^{1/2} \tilde{N}, \mathcal{D}^\circ \mathcal{D}^{1/4} \mathcal{D}, \mathcal{D}^3 \mathcal{D}^\circ \tilde{N} \cdot \tilde{N}, \tilde{N} \in \mathcal{D}^{3/4} \tilde{N} \cdot \mathcal{D}^\circ \mathcal{D}^{3/4} \mathcal{D} \zeta \mathcal{D}, \mathcal{D}, .$



Background From relatives and friends I heard about the gastroscopy and the horrors associated with this procedure.

Solutions I studied a number of sources â€” articles in the magazines â€œWhat doctors donâ€™t tell you aboutâ€•.

The option without anesthesia scared countless negative reviews.

Ingestion of the capsule is more suitable for colony, rather than gastroscopy, and there are only few centers and specialists in the procedure, high prices. And the need to go all day with the capsule inside and also with the device on the belt, then look for the capsule in your own waste and go to the doctor again. In addition, I read that the accuracy of this method leaves much to be desired. Gastroscopy in a dream is remained. The reviews are good. Nevertheless the need for anesthesia scared me. It took time to find a clinic and sign up for a procedure. By the way, the nearest record available was after about 2 weeks.

Preparation for gastroscopy in a dream By telephone medicine provider told me about the following: "A cardiogram with a decoding (≤ 1 month old) required" It should be no respiratory or chronic bronchopulmonary diseases "In the evening before the procedure, stop eating until 20.00" On the day of the procedure it is excluded eating and drinking, taking medications, using gum and candy and smoking "On the day of the procedure it is permissible to brush your teeth." Before the procedure you have to go through a conversation with the anesthetist

I complied with all the telephone recommendations and arrived at the medical center half an hour before the appointed time. I was asked to drink an "Espumisan" solution and was sent for an initial consultation with a gastroscopy specialist. The doctor inquired about my complaints, after which she sent me to the anesthesiologist. The anesthesiologist was interested in my chronic diseases, taking medications, studied my cardiogram, found out height, age and weight. He outlined the procedure for gastroscopy. Then I went to the procedure itself. I was asked to put on a robe and lie on the couch, on my left side. My lips were smeared with a moisturizer, and then was inserted a mouthpiece. Into a vein was inserted a catheter and was connected a pressure/pulse measurement. The first test dose of anesthesia was introduced to identify possible adverse reactions. I got dizzy. An iced solution was injected into the throat and the main dose of the drug was started. I have been slept instantly. I woke up on my own after 40 minutes. I was lying on the back, without a catheter and mouthpiece, under a blanket. I was informed that the gastroscopy lasted 7 minutes. Nausea and sensation in the stomach were not observed. It seemed that I slept for a maximum of half a minute, but at the same time I felt rested. I was asked to sit on the couch and wait for a couple of minutes, after which doctors helped me get up. After making sure that I was standing confidently, they asked me to go into the corridor and wait for the result of gastroscopy.

After about 10 minutes the doctor called me and reported the results. She gave me a color printout with characteristic moments of the gastroscopy. After that I sat in the corridor for about 20 minutes. Then I went to the clinic's cafe, where I was able to have breakfast without any restrictions.

